

# Mr Simon Grodski

## Postoperative Instructions

### HEMITHYROIDECTOMY PARTIAL THYROIDECTOMY



Mr Simon Grodski  
Suite 56  
Cabrini Hospital  
183 Wattletree Rd.  
Malvern, 3144

Ph. 9500-2288  
Fax. 9509-7109  
Mobile. 0402-434123  
e-mail. [info@simongrodski.com.au](mailto:info@simongrodski.com.au)

Suite 56, Cabrini Hospital  
183 Wattletree Rd.  
Malvern, 3144

Ph. 9500-2288

Fax. 9509-7109

## **HEMI THYROIDECTOMY PARTIAL THYROIDECTOMY**

### **Wound Care:**

Your wound will be covered with tape, which should be left in place for about 2 weeks. The tape will be removed at your first postoperative visit. You may notice some dried blood under the tape but that is of no concern. You can wash and shower with the tape in place and even get it wet. Once removed, you can shower with the wound exposed. The tape can then be replaced each day for 4-6 weeks after surgery in order to get the best possible cosmetic result. The preferred tape is ½ inch, flesh coloured Micropore tape which can be obtained from your chemist.

### **Activities:**

You should generally restrict vigorous activities for 1 or 2 weeks after surgery. Activities, which involve turning the head suddenly, such as driving in heavy traffic, should be avoided. Commonsense is the best way to avoid straining your neck.

### **Local symptoms:**

A variety of local symptoms are common for several weeks after surgery including tightness, swelling of the local area around the wound, choking, or having to force down food. Headaches and tiredness are also common as is weakness of the voice with prolonged use. Occasionally a symptom may last for

months. Swelling of the neck around the wound is also common and may benefit from daily massage with Vitamin E cream. Numbness of the skin above the wound may be present and may last for many months.

### **Late complications:**

The only delayed complication of concern is wound infection. This may have occurred if the wound becomes very red, hot and more swollen. If that occurs you must seek attention from your local doctor straight away who will arrange for you to have antibiotics.

### **Follow-up:**

Generally your follow-up will consist of

- A visit at 2 to 3 weeks to have the tape removed. You will need to ring the office to make an appointment for that visit.
- A final surgical check up at 3 months

In addition, you will also need to be reviewed by your local doctor or endocrinologist at around 6 weeks and have thyroid function tests to check the function of your remaining thyroid. Your local doctor or endocrinologist may need to see you more frequently for specific review if there are any problems.

### **Country patients:**

If you are unable to return to Melbourne for follow-up, it may be possible for the tape to be

changed and for your follow-up to be undertaken by your local doctor. We would need to confirm these arrangements before you leave hospital.

**If you have any questions or concerns please contact Mr Grodski as listed overleaf.**